



HHS Public Access

Author manuscript

Nat Rev Cancer. Author manuscript; available in PMC 2020 May 01.

Published in final edited form as:

Nat Rev Cancer. 2019 May ; 19(5): 271–282. doi:10.1038/s41568-019-0135-7.

Targeting cancer vulnerabilities with high-dose vitamin C

Bryan Ngo¹, Justin M. Van Riper², Lewis C. Cantley^{1,*}, and Jihye Yun^{2,*}

¹Meyer Cancer Center, Department of Medicine, Weill Cornell Medicine, New York, NY, USA.

²Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA.

Abstract

Over the past century, the notion that vitamin C can be used to treat cancer has generated much controversy. However, new knowledge regarding the pharmacokinetic properties of vitamin C and recent high-profile preclinical studies have revived interest in the utilization of high-dose vitamin C for cancer treatment. Studies have shown that pharmacological vitamin C targets many of the mechanisms that cancer cells utilize for their survival and growth. In this Opinion article, we discuss how vitamin C can target three vulnerabilities many cancer cells share: redox imbalance, epigenetic reprogramming and oxygen-sensing regulation. Although the mechanisms and predictive biomarkers that we discuss need to be validated in well-controlled clinical trials, these new discoveries regarding the anticancer properties of vitamin C are promising to help identify patient populations that may benefit the most from high-dose vitamin C therapy, developing effective combination strategies and improving the overall design of future vitamin C clinical trials for various types of cancer.

The ‘magic bullet’ theory serves as a paradigm for modern cancer research and has inspired numerous ground-breaking targeted therapies such as imatinib and vemurafenib¹. However, despite remarkable initial responses, the eventual acquisition of resistance and therapy-associated toxic effects continues to impede progress towards achieving meaningful patient survival. Thus, a new strategy for treating and managing cancer is needed. In this Opinion article, we propose that vitamin C, a natural compound with an unusually high safety profile, can be used to target multiple critical pathways in cancer.

The utilization of high-dose vitamin C as a cancer therapy has a controversial history. Much of this controversy stems from conflicting results in early clinical trials, as well as the lack of biomarkers and a clear understanding of vitamin C’s mechanism of action. Despite this,

* LCantley@med.Cornell.edu; Jihye.Yun@bcm.edu.

Author contributions

L.C.C. and J.Y. contributed to the discussion of content of the article. J.Y. and B.N. researched data for the article. J.Y. and B.N. wrote the article. All authors reviewed and edited the manuscript before submission.

Competing interests

The authors declare no competing interests.

Publisher’s note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Reviewer information

Nature Reviews Cancer thanks J. Cullen, G. Dachs and the other anonymous reviewer(s) for their contribution to the peer review of this work.

publications over the past 40 years suggest that these contradictory results can be explained, at least in part, by differences in the administration route of vitamin C; the millimolar concentration of vitamin C cytotoxic to cancer cells is achievable only by intravenous injection, not by oral administration². As a result, there are approximately a dozen ongoing clinical trials exploring the safety and efficacy of intravenous high-dose vitamin C for treating various types of cancer as a monotherapy or combination therapy³ (TABLE 1). Given the revived clinical interest in vitamin C as a cancer therapy, this Opinion article examines the evidence supporting the therapeutic potential of vitamin C and highlights advances in the current understanding of its mechanisms of action. First, we summarize the biological functions and chemical properties of vitamin C. Second, we examine three different mechanisms by which high-dose vitamin C can selectively kill cancer cells. Understanding the multiple targets and mechanisms by which vitamin C exerts anticancer effects will be essential for identifying predictive biomarkers for patient stratification and developing potent combination strategies that lead to durable remission. Finally, we close our Review by sharing our perspective on the future of vitamin C research as a treatment for cancer.

Biology of vitamin C

Synthesis of vitamin C

Vitamin C is a six-carbon ketolactone synthesized from glucose by most animals in the kidney or liver⁴. However, humans — as well as other primates, guinea pigs and fruit bats — are unable to synthesize vitamin C because they harbour inactivating mutations in the gene encoding L-gulonolactone oxidase (GULO), the enzyme responsible for catalysing the last step of vitamin C synthesis⁵. Owing to this ‘inborn metabolic error’, humans must acquire vitamin C from dietary sources. The current recommended daily allowance of vitamin C (75–90 mg per day) can easily be achieved by consuming a balanced diet consisting of fruits and vegetables, yielding a plasma ascorbate concentration of 30–80 μM (REF⁴). By contrast, sustained malnutrition or low dietary vitamin C intake will lead to plasma levels below 10 μM and result in scurvy, a vitamin C deficiency disease characterized by bleeding gums, impaired wound healing, anaemia, fatigue, depression and, in severe cases, death⁴.

Redox forms of vitamin C

Vitamin C exists in different redox forms depending on the biological conditions (FIG. 1a). Fully reduced vitamin C (ascorbate or ascorbic acid) can be oxidized both intracellularly and extracellularly. Extracellular ascorbate is oxidized by free radicals or reactive oxygen species (ROS) producing a weak radical intermediate, ascorbate radical ($\text{Asc}^{\bullet-}$), which is then oxidized fully into dehydroascorbic acid (DHA)⁶. DHA, having a short half-life (less than 1 minute)⁷, accounts for only approximately 1–5% of vitamin C in the human body⁴ and is either transported inside the cell (BOX 1) or becomes irreversibly hydrolysed into 2,3-L-diketoglutarate (2,3-DKG). 2,3-DKG is then degraded into oxalic acid and threonic acid, resulting in a net loss of vitamin C⁸. Inside the cell, DHA is rapidly reduced back to ascorbate by reacting with a reduced glutathione (GSH)⁸. Oxidized glutathione (glutathione disulfide (GSSG)) is then recycled back to GSH by NADPH⁸.

Biological functions of vitamin C

The biological functions of vitamin C can be attributed to its biochemical property as an electron donor. Acting as an antioxidant, physiological ascorbate at micromolar concentrations can reduce harmful ROS⁹. Paradoxically, it can also function as a pro-oxidant at millimolar plasma concentrations, which can be achieved by intravenous administration of pharmacological ascorbate⁴. In addition to its redox functions, vitamin C affects iron metabolism by increasing ferritin synthesis, inhibiting ferritin degradation, suppressing iron efflux and enhancing intestinal absorption of iron¹⁰. Interestingly, vitamin C also functions as a critical cofactor for numerous enzymes by readily donating its electrons to prosthetic metal ions to achieve full enzymatic activity⁴. In general, these enzymes are categorized into two families: copper-containing monooxygenases and Fe²⁺-dependent and α -ketoglutarate (α KG; also known as 2-oxoglutarate (2OG))-dependent dioxygenases (α KGDDs). α KGDDs are iron-containing enzymes that consume oxygen and α KG as co-substrates while producing CO₂ and succinate. α KGDDs catalyse a wide range of hydroxylation reactions involved in collagen synthesis, hypoxia-inducible factor 1 α (HIF1 α) stability, carnitine synthesis, the catabolism of tyrosine and the demethylation of protein, DNA and RNA. Thus, vitamin C is responsible for regulating a variety of important biological processes¹¹.

Anticancer mechanism of vitamin C

Over the past decade, a growing number of studies have demonstrated that millimolar concentrations of pharmacological vitamin C can kill cancer cells in vitro and slow tumour growth in vivo. However, the mechanism by which some cancer cells are sensitive to vitamin C, while normal cells remain resistant, is poorly understood. Given the diversity of processes affected by vitamin C, the mechanistic basis for vitamin C's action could depend on a variety of different factors, including the type of cancer being treated, and the tumour's dependency on particular pathways. Here, we discuss three distinct vulnerabilities in cancer that can be exploited by pharmacological ascorbate.

Targeting redox imbalance

It is generally accepted that cancer cells experience more oxidative stress compared with normal cells owing to an elevated metabolic rate and defective mitochondria¹². Although ROS can facilitate tumour development by stimulating cell proliferation and promoting genetic instability, excessive ROS can also be detrimental to cancer cells. To compensate, cancer cells often enhance pathways that help mitigate the toxic effects of ROS¹³. On the basis of the premise that ROS promotes cancer development, antioxidant treatment has been investigated as an anticancer strategy. However, recent results from both human and animal studies have found no clear evidence of the benefit of antioxidant treatment in preventing or suppressing cancer development. In some cases, antioxidant treatment even appeared to accelerate cancer progression and metastasis in mouse models of lung adenocarcinoma and melanoma^{14–16} and increase the risk of prostate and lung cancers in patients^{17–19}. Together, these results indicate that certain cancer types may rely on antioxidants for survival and may thus be vulnerable to pro-oxidant therapies. Indeed, pro-oxidant anticancer therapies, such as radiation, have been employed in the clinic²⁰. However, current pro-oxidant strategies often

cause serious collateral damage, resulting in a narrow therapeutic window²⁰. Here, we propose that pharmacological ascorbate can potentially circumvent this problem by exploring two common features of cancer cells: their increased levels of labile transition metals, especially iron²¹, and their increased reliance on glucose uptake and glycolysis²². Although we discuss these two mechanisms individually in this section, they are not mutually exclusive and can occur simultaneously, synergizing the selective toxicity of ascorbate in cancer cells (FIG. 1b).

Increased labile iron level.—In the presence of redox-active transition metals, such as iron, vitamin C exerts pro-oxidant effects. Iron is an essential prosthetic metal ion for a number of proteins²³. Most organisms require iron owing to its unique ability to efficiently switch between two oxidation states — ferrous iron (Fe^{2+} , reduced) and ferric iron (Fe^{3+} , oxidized) — in response to changes of ligands in the environment. Because of iron's highly reactive biochemical property, most labile iron must be sequestered by transferrin (Tf) in plasma, stored in ferritin inside the cell or embedded as cofactors, such as haem, in proteins²⁴. However, cells also keep small pools (~3–5% or ~1 μM in humans) of loosely coordinated Fe^{2+} , called labile iron pools, in the cytosol and the mitochondrial matrix for easy access²⁴. When labile Fe^{2+} in these pools reacts with H_2O_2 , it can generate the damaging hydroxyl radical ($\cdot\text{OH}$) via the Fenton reaction²¹ (FIG. 1b). To perpetuate this reaction, ascorbate effectively donates electrons to Fe^{3+} to regenerate redox-active Fe^{2+} , thereby generating ROS continuously and contributing to cell death²⁵. Thus, although iron is essential for a variety of biological processes, it can be a dangerous liability at the same time.

Numerous in vitro cell culture studies have shown that pharmacological ascorbate produces extracellular H_2O_2 , which can directly kill cancer cells^{26–28}. However, the exact mechanisms for this observation are currently unclear. Some studies showed that extracellular H_2O_2 can be generated via spontaneous autoxidation even in the absence of iron by reacting with oxygen when supraphysiological, millimolar concentrations of ascorbate are added to the medium in both cell-free and cell culture systems^{29,30}. Other studies demonstrated that labile metals, especially Fe^{2+} in the medium, catalyse ascorbate autoxidation, thereby generating extracellular H_2O_2 in both cell-free and cell culture systems^{31,32}. Because most labile Fe^{2+} is known to be sequestered by Tf in vivo²⁴, these studies argue that ascorbate's ability to produce H_2O_2 is an in vitro artefact^{33,34}. Disputing against this hypothesis, it was shown that $\text{Asc}^{\cdot-}$ and H_2O_2 were generated in vivo following intravenous ascorbate injections in rats (0.5 g kg^{-1}), and the production was ascorbate-dose-dependent³⁵. In another study, daily intraperitoneal injection with high-dose ascorbate (4 g kg^{-1}) inhibited neuroblastoma growth in a xenograft model, and tumours had the increased activity of checkpoint kinase 2 (CHK2) and histone 2AX (H2AX). This observation suggests that pharmacological ascorbate can cause DNA damage in in vivo tumours, although an in vivo link between ascorbate and the generation of H_2O_2 has not been demonstrated in this study³⁶.

The important question then is how extracellular H_2O_2 generated from pharmacological ascorbate can contribute to the selective toxicity to cancer cells compared with normal cells. If ascorbate generates extracellular H_2O_2 equally in both cancer cells and normal cells

regardless of the exact mechanism (for example, autoxidation and/or labile iron reactions in the medium or the serum in vivo), pharmacological ascorbate would not provide any additional advantage over other pro-oxidant therapies in terms of therapeutic window. Here, we propose three potential mechanisms by which ascorbate-induced H_2O_2 selectively kills certain types of cancer cells compared with normal cells. First, if the tumour microenvironment is enriched for labile iron, its reaction with ascorbate will generate H_2O_2 and $\cdot OH$, which can be lethal to cancer cells. Second, if tumour cells have increased levels of intracellular labile iron compared with normal cells, extracellular H_2O_2 from ascorbate autoxidation can diffuse into tumour cells and react with the intracellular pools of labile iron and generate $\cdot OH$ within tumour cells. Third, extracellular H_2O_2 can contribute to the increased level of extracellular DHA by generating an oxidative microenvironment. As a result, DHA is transported into tumours that express high levels of glucose transporter 1 (GLUT1), generating oxidative stress in those cells, which is discussed in the following section. In this section, we discuss the first and second possibilities.

Numerous iron-containing and haem-containing enzymes are involved in many important cellular processes such as cellular respiration, DNA synthesis, cell cycle and epigenetics^{23,36}. For this reason, cancer cells have a high demand for easily accessible labile Fe^{2+} for their survival and growth. It has been shown that reprogramming of iron metabolism can occur via a variety of mechanisms, including the upregulation of several iron-intake pathways or downregulation of iron export and storage pathways in various types of cancer such as breast, prostate and lymphoma^{37,38}. For example, the pool of intracellular labile Fe^{2+} in breast cancer cells is approximately twice as high as in normal breast epithelial cells³⁹. In addition, tumour-associated macrophages may promote iron release in the tumour microenvironments^{40,41}. Patients with advanced breast cancer have significantly higher levels of Fe^{2+} in plasma than healthy human control groups⁴².

Thus, tumour cells with the high levels of extracellular and/or intracellular labile iron may be more vulnerable to high-dose ascorbate than normal cells as they can generate more H_2O_2 and $\cdot OH$ than normal cells (FIG. 1b). Supporting this hypothesis, a recent study showed that an increased level of mitochondrial ROS in lung and glioblastoma cancer cells led to increased levels of intracellular labile iron through the upregulation of the Tf receptor (TfR), thus increasing the sensitivity of these cancer cells to ascorbate⁴³. Another study indicated that multiple myeloma cells have an elevated labile iron pool owing to the low expression of the iron exporter, ferroportin 1, which led to their selective sensitivity to pharmacological doses of ascorbate⁴⁴. Despite these promising preclinical studies, researchers still need to determine whether there is any correlation between ascorbate sensitivity, ascorbic radical or H_2O_2 production and intracellular labile Fe^{2+} levels in a large set of cancer cell lines or patient samples. Moreover, identifying potential biomarkers or gene expression signatures to predict ascorbate sensitivity related to the increased labile iron level would be useful in the clinic.

Increased DHA uptake via GLUT1.—Tumour cells exhibit a high rate of glycolysis, even in conditions with ample oxygen — a phenomenon that was first described by Otto Warburg, nearly a century ago²². This metabolic reprogramming, also known as the Warburg effect, is essential for tumour survival and proliferation⁴⁵. Oncogenic *KRAS* or *BRAF*

KRAS or *BRAF* mutant cells), they have an increased level of antioxidant enzymes and enhanced glucose flux into the pentose phosphate pathway (ppp)^{60,61} to generate more NADPH. In a normal setting, this protects erythrocytes from high-dose ascorbate. However, human erythrocytes without G6PD, the rate-limiting enzyme in PPP, cannot produce enough NADPH, a critical molecule for recovering depleted levels of GSH caused by vitamin-C-induced oxidative stress, which leads to the death of erythrocytes, thereby causing anaemia. For this reason, patients who plan to receive intravenous vitamin C therapy should be pre-screened for G6PD deficiency to avoid this complication. The fact that the most obvious and immediate oxidative stress effects of high-dose ascorbate in this genetic disorder occur in erythrocytes indicates the importance of increased GLUT1 levels and dependency on glycolysis in ascorbate-induced oxidative stress.

Targeting epigenetic regulators

Epigenetic reprogramming in cancer includes DNA hypermethylation, which frequently occurs on CpG island promoter regions and is known to silence tumour suppressors such as the retinoblastoma and VHL tumour suppressors⁶². Most of the aberrant DNA hypermethylation patterns observed in cancer can be explained by mutations or altered expression of two protein families: gain of function of DNA methyltransferases (DNMTs; which methylate cytosine to generate 5-methylcytosine (5mC)) and loss of function of ten-eleven translocation (TET) proteins.

TET proteins (TET1, TET2 and TET3) demethylate DNA and belong to the α KGDD enzyme family. Using oxygen and α KG, they catalyse multiple oxidation reactions, first converting 5mC to 5-hydroxymethylcytosine (5hmC), which ultimately results in an unmodified cytosine (FIG. 2). TET2 is frequently mutated or lost somatically in both myeloid and lymphoid malignancies^{63,64}. Interestingly, in a large acute myelogenous leukaemia (AML) cohort, TET2 was found to be mutually exclusive with gain-of-function mutations in isocitrate dehydrogenase 1 (IDH1) and IDH2 (REF.⁶⁵). IDH1 and IDH2 convert isocitrate to α KG in the cytosol and mitochondria, respectively. However, a neomorphic mutation in IDH1 and/or IDH2 causes a change in enzymatic activity and the accumulation of 2-hydroxyglutarate (2-HG), which inhibits the function of α KG-dependent enzymes such as TET2, resulting in the loss of 5hmC, an increase in DNA methylation and ultimately altered gene expression programmes that drive cancer development⁶⁶.

Vitamin C can activate TETs as a cofactor and is required for optimal activity. Ascorbate can donate an electron to Fe^{3+} to generate Fe^{2+} , which is required for TET activity (FIG. 2). The majority of TET2 mutations in AML are heterozygous, and each TET isoform has some functional redundancy. Therefore, ascorbate treatment may enhance the activity of residual TET proteins and thus rescue the abnormal DNA methylation pattern. Indeed, the growing number of recent studies supports this hypothesis. For example, daily intraperitoneal injection of high-dose ascorbate (4 g kg^{-1}) treatment in an inducible TET2 deletion mouse model of leukaemia recapitulated the TET2 restoration phenotypes by promoting DNA demethylation and the expression of genes critical for myeloid cell differentiation⁶⁷. Similarly, ascorbate treatment in vitro increased DNA demethylation at enhancers and promoters of genes associated with myeloid differentiation and increased the expression of

several key haematopoietic genes in murine bone marrow cells expressing mutant IDH1 (REF.⁶⁸). In certain types of lymphomas where TETs are frequently mutated, ascorbate treatment in vitro increased TET activity, leading to DNA demethylation, increased expression of tumour suppressor genes and increased chemosensitivity⁶⁹. Of note, all these studies applied appropriate controls to exclude oxidative stress in response to high-dose ascorbate as a possible mechanism by adding catalase (which converts H₂O₂ to water) to culture media, monitoring changes in cellular ROS levels and/or using 2-phosphate L-ascorbic acid, a vitamin C derivative that is stable and not oxidized in typical culture conditions, in their experiments. In addition to blood cancers, ascorbate treatment in melanoma and bladder cancer cells also enhanced 5hmC levels and decreased their malignancy^{70,71}, suggesting that ascorbate treatment may also be effective in solid cancer with low levels of 5hmC. Taken together, the results from these preclinical models warrant further investigation in the utilization of vitamin C therapy in patients with cancer who present with decreased levels of 5hmC and/or decreased TET activity.

A recent study also suggests that oral ascorbate may have a preventive role in leukaemogenesis⁷². Using several different genetically engineered mouse models (GEMMs) including *Gulo*^{-/-} mice, which are unable to produce vitamin C, and *Tet2*^{-/-} mice, ascorbate deficiency was shown to dysregulate HSC function in both a TET2-dependent and TET2-independent manner, leading to leukaemogenesis. On the basis of this study and the observational human studies (BOX 2), it would be important to determine the adequate amount of oral ascorbate to optimize human health and prevent chronic diseases such as cancer.

In addition to TETs, other αKGDDs such as JmjC domain-containing histone demethylase family (JHDM) and αKGDD AlkB (ALKB) are also known to be epigenetic regulators that can be potentially regulated by vitamin C^{73,74}. However, it is currently unclear whether these enzymes have important roles in cancer development and whether ascorbate availability influences their activity, thereby affecting the growth and survival of cancer cells.

Targeting HIF1 signalling

Many solid tumours encounter hypoxia as tumour masses can obstruct and compress surrounding blood vessels and tumour cells can outgrow new blood vessels. To adapt to this hypoxic microenvironment, tumour cells activate the evolutionarily conserved transcription factor HIF1, leading to activation of a wide range of genes and response programmes that facilitate increased survival in such conditions⁷⁵.

HIF1, an important target in cancer therapy, is a heterodimeric transcription factor, consisting of two subunits, the O₂-regulated HIF1α and a constitutively expressed HIF1β (REF.⁷⁵). The key mechanism by which O₂ regulates HIF1α activity is through proline hydroxylase domain proteins (PHD1, PHD2 and PHD3) and asparagine hydroxylase (factor-inhibiting HIF (FIH)), collectively known as the HIF hydroxylases. Under normoxic conditions, PHDs hydroxylate proline residues on HIF1α. Prolyl-hydroxylated HIF1α is then bound by the VHL tumour suppressor protein, which recruits an E3-ubiquitin ligase that targets HIF1α for proteasomal degradation (FIG. 3). On the other hand, FIHs

hydroxylate an asparagine residue on HIF1 α , which blocks the association of HIF1 α with the p300 co-activator protein, resulting in the inhibition of the transcription activity of HIF 1 (FIG. 3). Similar to TETs, both HIF hydroxylases belong to Fe²⁺-containing α KGDDs that require O₂ and α KG as substrates. Because of their relatively lower affinity for O₂ (Michaelis constant (K_m) = 230–250 μ M) compared with other α KGDDs⁷⁶, under hypoxic conditions, PHDs and FIHs are inactive, leading to the stabilization and activation of HIF1. Similar to TETs, HIF hydroxylases require ascorbate as a cofactor to recycle Fe²⁺. Therefore, cells deficient in ascorbate can have increased HIF1 α function, potentially contributing to tumour progression. This implies that ascorbate treatment may enhance the activity of HIF hydroxylases, thus inhibiting HIF1 α activity and suppressing tumour growth⁷⁷. Supporting this notion, there is growing evidence that HIF1 α -dependent tumour growth may be inhibited by ascorbate^{78,79}. Ascorbate level was inversely correlated with HIF1 α expression in thyroid lesions⁸⁰, and in vitro study showed that ascorbate treatment induced a dose-dependent decrease in expression of HIF1 α and GLUT1 (a downstream target of HIF 1) in thyroid cancer cells⁸⁰. In studies with *Gulo*^{-/-} mice, lung carcinoma implanted in *Gulo*^{-/-} mice grew slowly when mice were treated with high-dose ascorbate either in drinking water (3.3 g per litre) or daily intraperitoneal injection (1 gkg⁻¹) as compared with *Gulo*^{-/-} mice treated with low-dose ascorbate in drinking water (0.33 g per litre), and tumours exposed to high-dose ascorbate also had reduced expression levels of HIF1 α , VEGF and reduced microvessel density compared with control mice^{81–83}. In addition, retrospective human studies also support the connection between vitamin C, HIF1 activity and tumorigenesis. Using human patient tumour samples and paired controls for endometrial cancer, renal cell carcinoma (RCC) and CRC, it was demonstrated that tumours that had the highest HIF1 activity were those deficient in ascorbate in tumours^{84–86}. It was also shown that patients with CRC who had high levels of ascorbate in their tumour had a better patient outcome and longer patient survival after surgery⁸⁴. Taken together, these data suggest that high-dose ascorbate treatment can slow tumour growth by moderating HIF1 α . However, all studies thus far prove the association but not the causality. It would be interesting to see whether the deletion of PHDs or FIHs in tumour-bearing *Gulo*^{-/-} mice abolishes the effects of high-dose ascorbate.

The stabilization of HIF1 α even under normoxic conditions can occur, for example, in RCC, where deletion of the VHL tumour suppressor prevents HIF1 α degradation in normoxia. VHL-deficient RCC cells undergo cell death when exposed to vitamin C in normoxia, in contrast to isogenic VHL-proficient cells⁵². Mechanistically, the higher levels of GLUT1, a HIF1 α downstream target, in normoxic VHL-deficient cells than in VHL-proficient cells facilitated increased uptake of ascorbate-derived DHA, leading to increased generation of ROS and cell death. In line with this, GLUT1 knockdown in VHL-deficient RCC cells conferred resistance to vitamin-C-induced toxicity. In addition to VHL mutations, HIF1 α stability and activity can also be increased by mutations of two tumour suppressor enzymes in the tricarboxylic acid (TCA) cycle: succinate dehydrogenase (SDH) and fumarate dehydrogenase (FH). Loss-of-function mutations in SDH and FH cause a build-up of succinate and fumarate, respectively⁸⁷. Accordingly, increased levels of succinate and fumarate caused by mutations in these genes can compete with α KG and inhibit activity of HIF hydroxylases and thus induce normoxic HIF1 α activity in vitro^{88,89}. Because inherited

or somatic mutations in SDH and FH are tightly associated with the development of several tumours, such as paraganglioma, pheochromocytoma and RCC^{90–92}, it would be interesting to investigate whether ascorbate treatment of these mutant cells leads to inhibition of HIF1 activity and decreased malignancy in vivo.

Vitamin C as anticancer therapy

High-dose vitamin C should not be viewed as a ‘one-size-fits-all’ modality for cancer treatment. Understanding the critical differences between oral and intravenous ascorbate administration routes and knowing which patients to treat will be critical for clinical trial designs and the approval of new ascorbate therapies. Clinical trials in the 1970s involving patients with terminal cancer showed that intravenous high-dose ascorbate extended patient survival in response to ascorbate treatment^{93,94}, whereas large-scale randomized controlled trials (RCTs) in the 1980s failed to confirm these initial findings because ascorbate was given orally instead^{95,96}. As it turns out, the differences between oral and intravenous administration routes can affect the maximum achievable plasma concentration in patients. In the initial trials, ascorbate was administered both intravenously and orally and achieved a peak plasma concentration of 6 mM. However, in later trials, where the same dose of ascorbate was administered orally, a peak plasma concentration of less than 200 μ M was achieved⁹⁷. It is now widely accepted that the millimolar concentration of ascorbate needed to induce cytotoxicity in cancer cells can be achieved only when administered intravenously⁹⁷. For example, a phase I clinical study revealed that ascorbate concentrations could safely reach 25–30 mM with intravenous infusion of 100 g of vitamin C⁹⁸. In this study, plasma concentrations around 10 mM were sustained for at least 4 hours, which, on the basis of preclinical studies, is sufficient to slow the growth of cancer cells.

An increased understanding of the clinical pharmacokinetics of ascorbate has provided confidence in revisiting the clinical potential of ascorbate. Consequently, over the past decade, there have been an increased number of phase I/II clinical trials and case reports testing the safety and efficacy of high-dose ascorbate as a treatment for various cancer types such as ovarian, brain, prostate and lung cancers as a monotherapy or in combination with radiation and other conventional chemotherapies^{3,99,100}. In short, a significant number of clinical studies to date have indicated that intravenous high-dose ascorbate is well tolerated in patients with minimal toxicity, improves the quality of life for patients and has demonstrated synergistic therapeutic effects as well as reduced side effects when combined with radiation and standard chemotherapies. However, many of these studies were not designed as large-scale RCTs; thus, the efficacy of high-dose ascorbate therapy remains to be determined. Below, we discuss the future direction of clinical research into ascorbate as a cancer therapy.

From bench to bedside

In this Review, we propose that vitamin C serves as an excellent example of a natural compound that targets multiple vulnerable nodes to inhibit cancer growth. Although we identified three distinct mechanisms by which high-dose ascorbate can inhibit cancer growth, it is possible that each mechanism may not work independently. For example,

subpopulations of *KRAS* or *BRAF* mutant CRC cells that were not completely killed via the pro-oxidant mechanism in vivo — potentially owing to their distance from blood vessels, poor perfusion¹⁰¹ or resistance to ROS — may still be affected by ascorbate therapy via inhibition of HIF1 signalling and/or activation of TET enzymes. In fact, reports show that many CRCs have low expression levels of TETs and relatively high expression levels of HIF1 α , implying that pharmacological ascorbate may attack multiple nodes in tumours with minimum toxicity^{102,103}, making it an ideal ‘magic bullet’ for cancer therapy.

Understanding the mechanisms of action gives us critical information about the patient populations who may receive the most benefit from ascorbate therapy. On the basis of current preclinical studies, ascorbate may be more effective in patients with cancer with mutations in *KRAS*, *BRAF*, *TET2*, *IDH1* and/or *IDH2*, *VHL*, *SH* or *FDH*. Pending validation, these could be used as putative predictive biomarkers for ascorbate therapy in clinical trials. Multiple omics assays such as genomics, transcriptomics, proteomics and metabolomics can be performed using a patient's tumour biopsy samples not only to test existing hypotheses but also to generate hypotheses in an unbiased and comprehensive way. Furthermore, a patient's urine, blood and stool samples can also be analysed for omics assays and compared before and after receiving ascorbate treatment. After treatment with pharmacological ascorbate, correlations can be made between the response and disease progression to discover prospective biomarkers and better define the pharmacodynamics of ascorbate. Once clinicians confirm, or identify, promising biomarkers in these early trials, they will be better able to design and optimize a large-scale RCT for ascorbate therapy using an appropriate patient population. Poorly designed clinical trials in the past have stunted critical research on therapeutic efficacy of ascorbate as a cancer treatment. We must learn from our mistakes and design more thorough trials if we hope to reach a conclusion regarding the benefits of ascorbate in cancer therapy. Moreover, although not discussed extensively in this Review, it would be interesting to investigate the potential benefit of oral vitamin C supplements to prevent cancer (BOX 2).

Combination therapy

Although pharmacological ascorbate alone has been shown to reduce the tumour growth in many different mouse models of cancer, the clinical potential of ascorbate as an anticancer therapy may also lie in its combined use with other cancer therapies^{3,11}. Numerous preclinical and clinical studies indicate that parenteral injection of pharmacological ascorbate does not interfere with chemotherapy or radiotherapy and may even act synergistically^{43,104,105}. Moreover, because the mechanisms of action of vitamin C are becoming better defined, we can propose vitamin C combinations in a more rational, hypothesis-driven manner. For example, on the basis of ascorbate's known DNA demethylation effects via TET activation, a recent study found that vitamin C enhanced the effect of the DNMT inhibitor 5-aza-CdR (decitabine) by promoting the demethylation of human endogenous retroviruses and amplifying the expression of interferon-stimulated genes, leading to cancer cell death^{106,107}. Because decitabine is approved by the US Food and Drug Administration for the treatment of patients with myelodysplastic syndromes, where *TET2* is also frequently deleted¹⁰⁸, it would be interesting to test these combinations in clinical trials.

Combining vitamin C with immune checkpoint inhibitors and immunomodulatory agents, such as cytotoxic T lymphocyte-associated protein 4 (CTLA4) receptor and programmed death 1 (PD-1)–programmed death ligand 1 (PD-L1) blockers, may also lead to durable therapeutic responses against a broad spectrum of cancers. A compelling hypothesis is offered by the utilization and combination of high-dose vitamin C, which is known to enhance the function of both innate and adaptive immune cells^{109,110}. Tumours exist in a complex immune milieu that includes neutrophils, macrophages and lymphocytes. Although not much is known about how ascorbate is utilized by the various cells that make up the tumour microenvironment, studies have shown that phagocytes and lymphocytes have ascorbate concentrations 10–100 times greater than plasma⁴. Moreover, a wealth of knowledge is emerging that highlights the influence of ascorbate in inflammatory response and immune cell function^{110,111}, which suggests that ascorbate may have synergistic effects when combined with current immunotherapy.

Concluding remarks

Despite the unprecedented popularity of ascorbate among the public as an anecdotal ‘cure-all’ remedy, much of its biological functions and pharmacological activity have remained elusive. However, recent discoveries about the diverse biological functions of ascorbate, and its relevance to cancer therapy, generated exciting and promising hypotheses regarding the use of ascorbate in the treatment of cancer. We have discussed the critical role of ascorbate in the function of TETs, PHDs and FIHs, which are all α KGDDs. Given that more than forty α KGDDs exist¹¹², it is possible that other α KGDDs could contribute to the anticancer mechanisms of ascorbate. Discovering novel roles of ascorbate and the pathways it regulates will aid in the identification of molecules that can be targeted to sensitize tumours to ascorbate treatment and lead to the development of novel combination therapies.

To fully elucidate the biological functions of ascorbate and its relevance to cancer development, researchers will need to utilize better *in vivo* models that recapitulate the human condition. GEMMs are powerful tools for studying the pathogenesis of cancer and examining the systemic effects of vitamin C *in vivo*¹¹³. Recent studies utilizing GEMMs have enhanced our understanding of anticancer properties of ascorbate and reinforced our understanding of its mechanism of action as it pertains to cancer^{47,67,72}. Also, *Gulo*^{-/-} mice are an ideal candidate for vitamin C research as they, similar to humans, cannot synthesize vitamin C *de novo* in the liver. A significant number of human studies consistently show that ascorbate treatment improved quality of life for patients with cancer, and in combination therapy, ascorbate protected normal tissues from toxicity caused by chemotherapy¹¹⁴. These effects are likely caused by non-cell autonomous mechanisms, and if so, whole organisms or co-culture systems, rather than cancer cell lines themselves, would be crucial to discovering mechanisms of ascorbate’s indirect effects on cancer cells.

In conclusion, high-dose intravenous ascorbate represents a promising and inexpensive anticancer therapeutic option that should be further explored in clinical trials. Given its low toxicity and low financial cost, ascorbate could become an important weapon in our arsenal against cancer, either acting as a single agent with predictive biomarkers or used in combination as an adjuvant therapy. Although we are still waiting on a definitive answer for

the clinical benefits of ascorbate therapy in cancer, current preclinical and early phase I/II clinical trial results suggest that Linus Pauling's claims regarding the therapeutic benefits of vitamin C therapy in cancer may not be so outrageous after all.

Acknowledgements

The authors thank S. Wang in the Yun laboratory for helpful discussions and proofreading of the manuscript. The authors apologize for any references left uncited owing to space limitations. This work was supported by the US National Institutes of Health (NIH) grant R35 CA197588 (L.C.C.), Stand Up to Cancer–American Association for Cancer Research grant SU2C-AACR-DT22–17 (L.C.C.), the Cancer Prevention and Research Institute of Texas (CPRIT) grant RR170039 (J.Y.), the US National Cancer Institute (NCI) grant 1K22CA216036 (J.Y.), the US National Science Foundation Graduate Research Fellowship Program grant DGE1257284 (B.N.) and NCI grant 1F99CA234950–01 (B.N.). L.C.C. is a founder and member of the senior advisory boards of Agios Pharmaceuticals and Petra Pharmaceuticals, which are developing novel therapies for cancer. The L.C.C. laboratory also receives financial support from Petra Pharmaceuticals.

Glossary

2-Phosphate l-ascorbic acid

A derivative of ascorbate that is not oxidized in culture or serum but releases ascorbate once it is inside the cells via hydrolysis mediated by alkaline phosphatase on the plasma membrane.

5-Aza-CdR (decitabine)

A cytidine antimetabolite analogue that incorporates into DNA and inhibits DNA methyltransferase (DNMT) activity which results in DNA demethylation (hypomethylation).

Biomarkers

Any biological measurable indicators of the severity or presence of some disease state.

Fenton reaction

A chemical reaction that converts hydrogen peroxide into a highly toxic hydroxyl radical in the presence of labile iron.

Ferritin

A protein that contains iron and is the primary form of iron stored inside of cells.

Free radicals

Molecules possessing unpaired electrons and thus are reactive and short-lived in a biological setting.

Haem

An iron-containing group that gives myoglobin and haemoglobin the ability to bind oxygen.

Hydroxyl radical

($\cdot\text{OH}$). A highly reactive and short-lived radical that attacks any molecule in its immediate vicinity, especially DNA, protein and lipids, eventually leading to cell death.

Imatinib

A BCR-ABL-selective tyrosine kinase inhibitor, also known as Gleevec. Imatinib has been used to treat chronic myelogenous leukaemia and acute lymphocytic leukaemia.

Intraperitoneal (IP) injection

Giving medicines or fluids into the peritoneum (body cavity), which is more often applied to animals than to humans.

Intravenous injection

Giving medicines or fluids through a needle or tube inserted into a vein, allowing them to enter the bloodstream immediately.

Michaelis constant

(K_m). The substrate concentration at the half of the maximum velocity (V_{max}). An enzyme with a high K_m has a low affinity for its substrate and requires a greater concentration of substrate to achieve V_{max} .

Parenteral injection

Giving medicine or fluids intravenously (into a vein), subcutaneously (under the skin) and intraperitoneally (into the peritoneum).

Pharmacodynamics

The study of the biochemical and physiological effects of drugs. Generally refers to the dose–response relationship for a particular drug.

Pharmacokinetics

The activity of drugs in the body over a period, including the processes by which drugs are absorbed, distributed in the body, localized in the tissues and excreted.

Pharmacological ascorbate

Intravenous or intraperitoneal delivery of vitamin C, which allows for plasma concentrations to reach the millimolar scale.

Physiological ascorbate

An oral dose of dietary vitamin C, usually resulting in a peak plasma concentration of 200 μM .

Predictive biomarkers

A biomarker that gives information about the effect of a therapeutic intervention.

Prosthetic

A group that is a tightly bound, specific non-polypeptide unit required for the biological function of some proteins. It may be organic or inorganic (such as a metal ion), but not amino acids.

Randomized controlled trials

(RCTs). A study design that randomly assigns participants into an experimental group or a control group (or placebo group).

Reactive oxygen species

(ROS). Derivatives of oxygen that are more reactive than molecular oxygen.

Single-nucleotide polymorphisms

(SNPs). A variation in a single nucleotide that occurs at a specific position in the genome, where each variation is present to some high degree within a population (for example, >1%).

Therapeutic window

The range of doses of a drug that can treat disease effectively without having toxic effects.

Transferrin

(Tf). The main protein in the blood that binds to iron and transports it throughout the body.

Vemurafenib

A selective V600E mutant BRAF kinase inhibitor, also known as PLX4032. It has been used to treat BRAF V600E mutant melanoma.

References

1. Strebhardt K & Ullrich A Paul Ehrlich's magic bullet concept: 100 years of progress. *Nat. Rev. Cancer* 8, 473–480 (2008). [PubMed: 18469827]
2. Levine M, Padayatty SJ & Espesy MG Vitamin C: a concentration-function approach yields pharmacology and therapeutic discoveries. *Adv. Nutr* 2, 78–88 (2011). [PubMed: 22332036]
3. Nauman G et al. Systematic review of intravenous ascorbate in cancer clinical trials. *Antioxidants* 7, E89 (2018). [PubMed: 30002308]
4. Padayatty SJ & Levine M Vitamin C: the known and the unknown and Goldilocks. *Oral Dis* 22, 463–493 (2016). [PubMed: 26808119]
5. Maeda N et al. Aortic wall damage in mice unable to synthesize ascorbic acid. *Proc. Natl Acad. Sci. USA* 97, 841–846 (2000). [PubMed: 10639167]
6. Corti A, Casini AF & Pompella A Cellular pathways for transport and efflux of ascorbate and dehydroascorbate. *Arch. Biochem. Biophys* 500, 107–115 (2010). [PubMed: 20494648]
7. Vera JC et al. Human HL-60 myeloid leukemia cells transport dehydroascorbic acid via the glucose transporters and accumulate reduced ascorbic acid. *Blood* 84, 1628–1634 (1994). [PubMed: 8068952]
8. Wilson JX The physiological role of dehydroascorbic acid. *FEBS Lett* 527, 5–9 (2002). [PubMed: 12220624]
9. Buettner GR The pecking order of free radicals and antioxidants: lipid peroxidation, alpha-tocopherol, and ascorbate. *Arch. Biochem. Biophys* 300, 535–543 (1993). [PubMed: 8434935]
10. Lane DJR & Richardson DR The active role of vitamin C in mammalian iron metabolism: much more than just enhanced iron absorption! *Free Radic. Biol. Med* 75, 69–83 (2014). [PubMed: 25048971]
11. Kuiper C & Vissers MCM Ascorbate as a co-factor for Fe- and 2-oxoglutarate dependent dioxygenases: physiological activity in tumor growth and progression. *Front. Oncol* 4, 359 (2014). [PubMed: 25540771]
12. Schieber M & Chandel NS ROS function in redox signaling and oxidative stress. *Curr. Biol* 24, R453–R462 (2014). [PubMed: 24845678]
13. Chio IIC & Tuveson DA ROS in cancer: the burning question. *Trends Mol. Med* 23, 411–429 (2017). [PubMed: 28427863]
14. Sayin VI et al. Antioxidants accelerate lung cancer progression in mice. *Sci. Transl Med* 6, 221ra15 (2014). This study demonstrates that dietary supplementation of antioxidants (N-acetylcysteine and vitamin E) markedly increases tumour progression and reduces survival in mouse models of KRAS-induced or BRAF-induced lung cancer.
15. Le Gal K et al. Antioxidants can increase melanoma metastasis in mice. *Sci. Transl Med* 7, 308re8 (2015).

16. Piskounova E et al. Oxidative stress inhibits distant metastasis by human melanoma cells. *Nature* 527, 186–191 (2015). [PubMed: 26466563]
17. Klein EA et al. Vitamin E and the risk of prostate cancer: the selenium and vitamin E cancer prevention trial (SELECT). *JAMA* 306, 1549–1556 (2011). [PubMed: 21990298]
18. Omenn GS et al. Effects of a combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *N. Engl. J. Med* 334, 1150–1155 (1996). [PubMed: 8602180]
19. Alpha-Tocopherol, Beta Carotene Cancer Prevention Study Group. The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. *N. Engl. J. Med* 330, 1029–1035 (1994). [PubMed: 8127329]
20. Wondrak GT Redox-directed cancer therapeutics: molecular mechanisms and opportunities. *Antioxid. Redox Signal* 11, 3013–3069 (2009). [PubMed: 19496700]
21. Torti SV & Torti FM Iron and cancer: more ore to be mined. *Nat. Rev. Cancer* 13, 342–355 (2013). [PubMed: 23594855]
22. DeBerardinis RJ & Chandel NS Fundamentals of cancer metabolism. *Sci. Adv* 2, e1600200 (2016). [PubMed: 27386546]
23. Manz DH et al. Iron and cancer: recent insights. *Ann. NY Acad. Sci* 1368, 149–161 (2016). [PubMed: 26890363]
24. Hentze MW et al. Two to tango: regulation of mammalian iron metabolism. *Cell* 142, 24–38 (2010). [PubMed: 20603012]
25. Du J, Cullen JJ & Buettner GR Ascorbic acid: chemistry, biology and the treatment of cancer. *Biochim. Biophys. Acta* 1826, 443–457 (2012). [PubMed: 22728050]
26. Chen Q et al. Pharmacologic ascorbic acid concentrations selectively kill cancer cells: action as a pro-drug to deliver hydrogen peroxide to tissues. *Proc. Natl Acad. Sci. USA* 102, 13604–13609 (2005). [PubMed: 16157892]
27. Du J et al. Mechanisms of ascorbate-induced cytotoxicity in pancreatic cancer. *Clin. Cancer Res* 16, 509–520 (2010). [PubMed: 20068072]
28. Rawal M et al. Manganoporphyrins increase ascorbate-induced cytotoxicity by enhancing H₂O₂ generation. *Cancer Res* 73, 5232–5241 (2013). [PubMed: 23764544]
29. Duarte TL, Almeida GM & Jones GDD Investigation of the role of extracellular H₂O₂ and transition metal ions in the genotoxic action of ascorbic acid in cell culture models. *Toxicol. Lett* 170, 57–65 (2007). [PubMed: 17382497]
30. Sakagami H et al. Effect on an iron-chelator on ascorbate-induced cytotoxicity. *Free Radic. Biol. Med* 23, 260–270 (1997). [PubMed: 9199888]
31. Buettner GR & Jurkiewicz BA Catalytic metals, ascorbate and free radicals: combinations to avoid. *Radiat. Res* 145, 532–541 (1996). [PubMed: 8619018]
32. Clement MV et al. The in vitro cytotoxicity of ascorbate depends on the culture medium used to perform the assay and involves hydrogen peroxide. *Antioxid. Redox Signal* 3, 157–163 (2001). [PubMed: 11291594]
33. Carr A & Frei B Does vitamin C act as a pro-oxidant under physiological conditions? *FASEB J* 13, 1007–1024 (1999). [PubMed: 10336883]
34. Halliwell B Vitamin C: antioxidant or pro-oxidant in vivo? *Free Radic. Res* 25, 439–454 (1996). [PubMed: 8902542]
35. Chen Q et al. Ascorbate in pharmacologic concentrations selectively generates ascorbate radical and hydrogen peroxide in extracellular fluid in vivo. *Proc. Natl Acad. Sci. USA* 104, 8749–8754 (2007). [PubMed: 17502596] This study shows that pharmacological ascorbate is a prodrug for preferential steady-state formation of Asc^{•-} and H₂O₂ in the extracellular space but not in the blood in vivo system.
36. Ma E et al. Pharmacologic ascorbate induces neuroblastoma cell death by hydrogen peroxide mediated DNA damage and reduction in cancer cell glycolysis. *Free Radic. Biol. Med* 113, 36–47 (2017). [PubMed: 28916476]
37. Rychtarcikova Z et al. Tumor-initiating cells of breast and prostate origin show alterations in the expression of genes related to iron metabolism. *Oncotarget* 8, 6376–6398 (2017). [PubMed: 28031527]

38. Kiessling MK et al. Inhibition of constitutively activated nuclear factor-kappaB induces reactive oxygen species- and iron-dependent cell death in cutaneous T cell lymphoma. *Cancer Res* 69, 2365–2374 (2009). [PubMed: 19258503]
39. Pinnix ZK et al. Ferroportin and iron regulation in breast cancer progression and prognosis. *Sci. Transl Med* 2, 43ra56 (2010).
40. Corna G et al. Polarization dictates iron handling by inflammatory and alternatively activated macrophages. *Haematologica* 95, 1814–1822 (2010). [PubMed: 20511666]
41. Recalcati S et al. Differential regulation of iron homeostasis during human macrophage polarized activation. *Eur. J. Immunol* 40, 824–835 (2010). [PubMed: 20039303]
42. Panis C et al. Differential oxidative status and immune characterization of the early and advanced stages of human breast cancer. *Breast Cancer Res. Treat* 133, 881–888 (2012). [PubMed: 22048816]
43. Schoenfeld JD et al. $O_2^{\bullet -}$ and H_2O_2 mediated disruption of Fe metabolism causes the differential susceptibility of NSCLC and GBM cancer cells to pharmacological ascorbate. *Cancer Cell* 31, 487–500. (2017). [PubMed: 28366679] This study shows that pharmacological ascorbate selectively sensitizes non-small-cell lung cancer (NSCLC) and glioblastoma (GBM) cells by generating redox-active labile iron and H_2O_2 using in vitro cell lines and mouse models. In addition, it shows the potential efficacy of ascorbate therapy combined with other chemotherapies in patients with GBM and NSCLC.
44. Xia J et al. Multiple myeloma tumor cells are selectively killed by pharmacologically-dosed ascorbic acid. *EBioMedicine* 18, 41–49 (2017). [PubMed: 28229908]
45. Liberti MV & Locasale JW The Warburg effect: how does it benefit cancer cells? *Trends Biochem. Sci* 41, 211–218 (2016). [PubMed: 26778478]
46. Yun J et al. Glucose deprivation contributes to the development of KRAS pathway mutations in tumor cells. *Science* 325, 1555–1559 (2009). [PubMed: 19661383]
47. Yun J et al. Vitamin C selectively kills KRAS and BRAF mutant colorectal cancer cells by targeting GAPDH. *Science* 350, 1391–1396 (2015). [PubMed: 26541605] This study demonstrates that uptake of DHA by GLUT1, and its subsequent reduction to ascorbate, depletes essential antioxidants to induce cell death in KRAS or BRAF mutant CRC cells.
48. Nualart FJ et al. Recycling of vitamin C by a bystander effect. *J. Biol. Chem* 278, 10128–10133 (2003). [PubMed: 12435736]
49. May JM et al. Ascorbate recycling in human erythrocytes: role of GSH in reducing dehydroascorbate. *Free Radic. Biol. Med* 20, 543–551 (1996). [PubMed: 8904295]
50. Tu H et al. Chemical transport knockout for oxidized vitamin C, dehydroascorbic acid, reveals its functions in vivo. *EBioMedicine* 23, 125–135 (2017). [PubMed: 28851583]
51. Lu Y-X et al. Pharmacological ascorbate suppresses growth of gastric cancer cells with GLUT1 overexpression and enhances the efficacy of oxaliplatin through redox modulation. *Theranostics* 8, 1312–1326 (2018). [PubMed: 29507622]
52. Tian W et al. The hypoxia-inducible factor renders cancer cells more sensitive to vitamin C-induced toxicity. *J. Biol. Chem* 289, 3339–3351 (2014). [PubMed: 24371136]
53. Vera JC et al. Resolution of the facilitated transport of dehydroascorbic acid from its intracellular accumulation as ascorbic acid. *J. Biol. Chem* 270, 23706–23712 (1995). [PubMed: 7559541]
54. Szatrowski TP & Nathan CF Production of large amounts of hydrogen peroxide by human tumor cells. *Cancer Res* 51, 794–798 (1991). [PubMed: 1846317]
55. Policastro LL et al. The tumor microenvironment: characterization, redox considerations, and novel approaches for reactive oxygen species-targeted gene therapy. *Antioxid. Redox Signal* 19, 854–895 (2013). [PubMed: 22794113]
56. Rees DC, Kelsey H & Richards JD Acute haemolysis induced by high dose ascorbic acid in glucose-6-phosphate dehydrogenase deficiency. *BMJ* 306, 841–842 (1993). [PubMed: 8490379]
57. Quinn J et al. Effect of high-dose vitamin C infusion in a glucose-6-phosphate dehydrogenase-deficient patient. *Case Rep. Med* 2017, 5202606 (2017). [PubMed: 29317868]
58. Rakitzis ET & Papandreou PT Ascorbate-induced generation of free radical species in normal and glucose-6-phosphate dehydrogenase-deficient erythrocytes. *Biochem. Soc. Trans* 17, 371–372 (1989).

59. Mehta J, Singhal S & Mehta B Ascorbic-acid-induced haemolysis in G-6-PD deficiency. *Lancet* 336, 944 (1990).
60. Gaetani GF et al. Predominant role of catalase in the disposal of hydrogen peroxide within human erythrocytes. *Blood* 87, 1595–1599 (1996). [PubMed: 8608252]
61. May JM, Qu Z & Morrow JD Mechanisms of ascorbic acid recycling in human erythrocytes. *Biochim. Biophys. Acta* 1528, 159–166 (2001). [PubMed: 11687303]
62. Yun J et al. Interactions between epigenetics and metabolism in cancers. *Front. Oncol* 2, 163 (2012). [PubMed: 23162793]
63. Rasmussen KD & Helin K Role of TET enzymes in DNA methylation, development, and cancer. *Genes Dev* 30, 733–750 (2016). [PubMed: 27036965]
64. Delhommeau F et al. Mutation in TET2 in myeloid cancers. *N. Engl. J. Med* 360, 2289–2301 (2009). [PubMed: 19474426]
65. Figueroa ME et al. Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell* 18, 553–567 (2010). [PubMed: 21130701]
66. Lu C et al. IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature* 483, 474–478 (2012). [PubMed: 22343901]
67. Cimmino L et al. Restoration of TET2 function blocks aberrant self-renewal and leukemia progression. *Cell* 170, 1079–1095 (2017). [PubMed: 28823558] Using genetic mouse models of leukaemia, vitamin C is shown to enhance TET2 activity, increase DNA demethylation and suppress leukaemia progression.
68. Mingay M et al. Vitamin C-induced epigenomic remodelling in IDH1 mutant acute myeloid leukaemia. *Leukemia* 32, 11–20 (2018). [PubMed: 28663574]
69. Shenoy N et al. Upregulation of TET activity with ascorbic acid induces epigenetic modulation of lymphoma cells. *Blood Cancer J* 7, e587 (2017). [PubMed: 28731456]
70. Gustafson CB et al. Epigenetic reprogramming of melanoma cells by vitamin C treatment. *Clin. Epigenetics* 7, 51 (2015). [PubMed: 25977731]
71. Peng D et al. Vitamin C increases 5-hydroxymethylcytosine level and inhibits the growth of bladder cancer. *Clin. Epigenetics* 10, 94 (2018). [PubMed: 30005692]
72. Agathocleous M et al. Ascorbate regulates haematopoietic stem cell function and leukaemogenesis. *Nature* 549, 476–481 (2017). [PubMed: 28825709] Using various GEMMs, including GULO-knockout mice, this study reveals that vitamin C is necessary for normal haematopoietic stem cell development and the prevention of leukaemia in mice.
73. Monfort A & Wutz A Breathing-in epigenetic change with vitamin C. *EMBO Rep* 14, 337–346 (2013). [PubMed: 23492828]
74. Fedeles BI et al. The AlkB family of Fe(II)/ α -ketoglutarate-dependent dioxygenases: repairing nucleic acid alkylation damage and beyond. *J. Biol. Chem* 290, 20734–20742 (2015). [PubMed: 26152727]
75. Semenza GL Targeting HIF-1 for cancer therapy. *Nat. Rev. Cancer* 3, 721–732 (2003). [PubMed: 13130303]
76. Koivunen P et al. Catalytic properties of the asparaginyl hydroxylase (FIH) in the oxygen sensing pathway are distinct from those of its prolyl 4-hydroxylases. *J. Biol. Chem* 279, 9899–9904 (2004). [PubMed: 14701857]
77. Vissers MCM & Das AB Potential mechanisms of action for vitamin C in cancer: reviewing the evidence. *Front. Physiol* 9, 809 (2018). [PubMed: 30018566]
78. Wilkes JG et al. Pharmacologic ascorbate (P-AscH-) suppresses hypoxia-inducible Factor-1 α (HIF-1 α) in pancreatic adenocarcinoma. *Clin. Exp. Metastasis* 35, 37–51 (2018). [PubMed: 29396728]
79. Gao P et al. HIF-dependent anti-tumorigenic effect of anti-oxidants in vivo. *Cancer Cell* 12, 230–238 (2007). [PubMed: 17785204]
80. Józwiak P et al. Expression of hypoxia inducible factor 1 α and 2 α and its association with vitamin C level in thyroid lesions. *J. Biomed. Sci* 24, 83 (2017). [PubMed: 29084538]
81. Campbell EJ et al. Restoring physiological levels of ascorbate slows tumor growth and moderates HIF-1 pathway activity in Gulo(-/-) mice. *Cancer Med* 4, 303–314 (2015). [PubMed: 25354695]

82. Campbell EJ, Vissers MC & Dachs GU Ascorbate availability affects tumor implantation-take rate and increases tumor rejection in Gulo^{-/-} mice. *Hypoxia* 4, 41–52 (2016). [PubMed: 27800507]
83. Campbell EJ et al. Pharmacokinetic and anti-cancer properties of high dose ascorbate in solid tumours of ascorbate-dependent mice. *Free Radic. Biol. Med* 99, 451–462 (2016). [PubMed: 27567539]
84. Kuiper C et al. Increased tumor ascorbate is associated with extended disease-free survival and decreased hypoxia-inducible factor-1 activation in human colorectal cancer. *Front. Oncol* 4, 10 (2014). [PubMed: 24551593]
85. Wohlrab C et al. The association between ascorbate and the hypoxia-inducible factors in human renal cell carcinoma requires a functional Von Hippel-Lindau protein. *Front. Oncol* 8, 574 (2018). [PubMed: 30555801]
86. Kuiper C et al. Low ascorbate levels are associated with increased hypoxia-inducible factor-1 activity and an aggressive tumor phenotype in endometrial cancer. *Cancer Res* 70, 5749–5758 (2010). [PubMed: 20570889]
87. King A, Selak MA & Gottlieb E Succinate dehydrogenase and fumarate hydratase: linking mitochondrial dysfunction and cancer. *Oncogene* 25, 4675–4682 (2006). [PubMed: 16892081]
88. Selak MA et al. Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF- α prolyl hydroxylase. *Cancer Cell*, 7, 77–85 (2005). [PubMed: 15652751]
89. Koivunen P et al. Inhibition of hypoxia-inducible factor (HIF) hydroxylases by citric acid cycle intermediates: possible links between cell metabolism and stabilization of HIF. *J. Biol. Chem* 282, 4524–4532 (2007). [PubMed: 17182618]
90. Gill AJ et al. Renal tumors and hereditary pheochromocytoma-paranglioma syndrome type 4. *N. Engl. J. Med* 364, 885–886 (2011). [PubMed: 21366490]
91. Pasini B & Stratakis CA SDH mutations in tumorigenesis and inherited endocrine tumours: lesson from the pheochromocytoma-paranglioma syndromes. *J. Intern. Med* 266, 19–42 (2009). [PubMed: 19522823]
92. Clark GR et al. Germline FH mutations presenting with pheochromocytoma. *J. Clin. Endocrinol. Metab* 99, E2046–E2050 (2014). [PubMed: 25004247]
93. Cameron E & Pauling L Supplemental ascorbate in the supportive treatment of cancer: prolongation of survival times in terminal human cancer. *Proc. Natl Acad. Sci. USA* 73, 3685–3689 (1976). [PubMed: 1068480] This study is the first large-scale clinical trial designed to analyse the efficacy of high-dose ascorbate for the treatment of cancer.
94. Cameron E & Pauling L Supplemental ascorbate in the supportive treatment of cancer: reevaluation of prolongation of survival times in terminal human cancer. *Proc. Natl Acad. Sci. USA* 75, 4538–4542 (1978). [PubMed: 279931]
95. Creagan ET et al. Failure of high-dose vitamin C (ascorbic acid) therapy to benefit patients with advanced cancer. A controlled trial. *N. Engl. J. Med* 301, 687–690 (1979). [PubMed: 384241] This large-scale clinical trial shows that daily oral intake of high-dose vitamin C had no therapeutic benefit in patients with cancer.
96. Moertel CG et al. High-dose vitamin C versus placebo in the treatment of patients with advanced cancer who have had no prior chemotherapy. A randomized double-blind comparison. *N. Engl. J. Med* 312, 137–141 (1985). [PubMed: 3880867]
97. Padayatty SJ et al. Vitamin C pharmacokinetics: implications for oral and intravenous use. *Ann. Intern. Med* 140, 533–537 (2004). [PubMed: 15068981] This paper emphasizes the importance of the route of administration of vitamin C and explains why two early clinical trials of vitamin C as an anticancer therapy had conflicting results.
98. Hoffer LJ et al. Phase I clinical trial of i.v. ascorbic acid in advanced malignancy. *Ann. Oncol* 19, 1969–1974 (2008). [PubMed: 18544557]
99. Klimant E et al. Intravenous vitamin C in the supportive care of cancer patients: a review and rational approach. *Curr. Oncol* 25, 139–148 (2018). [PubMed: 29719430]
100. Carr AC & Cook J Intravenous vitamin C for cancer therapy — identifying the current gaps in our knowledge. *Front. Physiol* 9, 1182 (2018). [PubMed: 30190680]
101. Kuiper C, Vissers MCM & Hicks KO Pharmacokinetic modeling of ascorbate diffusion through normal and tumor tissue. *Free Radic. Biol. Med* 77, 340–352 (2014). [PubMed: 25277418]

102. Cao D et al. Expression of HIF-1 α and VEGF in colorectal cancer: association with clinical outcomes and prognostic implications. *BMC Cancer* 9, 432 (2009). [PubMed: 20003271]
103. Tian Y et al. Association of TET1 expression with colorectal cancer progression. *Scand. J. Gastroenterol* 52, 312–320 (2017). [PubMed: 27846738]
104. Mustafi S et al. Vitamin C sensitizes melanoma to BET inhibitors. *Cancer Res* 78, 572–583 (2018). [PubMed: 29180474]
105. Du J et al. Pharmacological ascorbate radiosensitizes pancreatic cancer. *Cancer Res* 75, 3314–3326 (2015). [PubMed: 26081808]
106. Liu M et al. Vitamin C increases viral mimicry induced by 5-aza-2'-deoxycytidine. *Proc. Natl Acad. Sci. USA* 113, 10238–10244 (2016). [PubMed: 27573823]
107. Zhao H et al. The synergy of Vitamin C with decitabine activates TET2 in leukemic cells and significantly improves overall survival in elderly patients with acute myeloid leukemia. *Leuk. Res* 66, 1–7 (2018). [PubMed: 29331774]
108. Gillberg L et al. Vitamin C — a new player in regulation of the cancer epigenome. *Semin. Cancer Biol* 51, 59–67 (2017). [PubMed: 29102482]
109. Carr AC & Maggini S Vitamin C and immune function. *Nutrients* 9, 1211 (2017).
110. Sorice A et al. Ascorbic acid: its role in immune system and chronic inflammation diseases. *Mini Rev. Med. Chem* 14, 444–452 (2014). [PubMed: 24766384]
111. Ang A et al. Vitamin C and immune cell function in inflammation and cancer. *Biochem. Soc. Trans* 46, 1147–1159 (2018). [PubMed: 30301842]
112. Hausinger RP Fell/alpha-ketoglutarate-dependent hydroxylases and related enzymes. *Crit. Rev. Biochem. Mol. Biol* 39, 21–68 (2004). [PubMed: 15121720]
113. Kersten K et al. Genetically engineered mouse models in oncology research and cancer medicine. *EMBO Mol. Med* 9, 137–153 (2017). [PubMed: 28028012]
114. Ma Y et al. High-dose parenteral ascorbate enhanced chemosensitivity of ovarian cancer and reduced toxicity of chemotherapy. *Sci. Transl Med* 6, 222ra18 (2014). This paper shows in its preclinical and clinical studies that high-dose vitamin C can potentially increase the efficacy and reduce the toxicities of certain conventional chemotherapies when used in combination to treat patients with ovarian cancer.
115. May JM The SLC23 family of ascorbate transporters: ensuring that you get and keep your daily dose of vitamin C. *Br. J. Pharmacol* 164, 1793–1801 (2011). [PubMed: 21418192]
116. Corpe CP et al. Vitamin C transporter Slc23a1 links renal reabsorption, vitamin C tissue accumulation, and perinatal survival in mice. *J. Clin. Invest* 120, 1069–1083 (2010). [PubMed: 20200446]
117. Sotiriou S et al. Ascorbic-acid transporter Slc23a1 is essential for vitamin C transport into the brain and for perinatal survival. *Nat. Med* 8, 514–517 (2002). [PubMed: 11984597]
118. Vera JC et al. Mammalian facilitative hexose transporters mediate the transport of dehydroascorbic acid. *Nature* 364, 79–82 (1993). [PubMed: 8316303]
119. Spielholz C et al. Increased facilitated transport of dehydroascorbic acid without changes in sodium-dependent ascorbate transport in human melanoma cells. *Cancer Res* 57, 2529–2537 (1997). [PubMed: 9192836]
120. Agus DB, Vera JC & Golde DW Stromal cell oxidation: a mechanism by which tumors obtain vitamin C. *Cancer Res* 59, 4555–4558 (1999). [PubMed: 10493506]
121. Mayland CR, Bennett MI & Allan K Vitamin C deficiency in cancer patients. *Palliat. Med* 19, 17–20 (2005). [PubMed: 15690864]
122. Anthony HM & Schorah CJ Severe hypovitaminosis C in lung-cancer patients: the utilization of vitamin C in surgical repair and lymphocyte-related host resistance. *Br. J. Cancer* 46, 354–367 (1982). [PubMed: 7126425]
123. Ullah MF et al. Ascorbic acid in cancer chemoprevention: translational perspectives and efficacy. *Curr. Drug Targets* 13, 1757–1771 (2012). [PubMed: 23140287]
124. Lee KW et al. Vitamin C and cancer chemoprevention: reappraisal. *Am. J. Clin. Nutr* 78, 1074–1078 (2003). [PubMed: 14668266]

125. Khaw KT et al. Relation between plasma ascorbic acid and mortality in men and women in EPIC-Norfolk prospective study: a prospective population study. *European Prospective Investigation into Cancer and Nutrition. Lancet* 357, 657–663 (2001). [PubMed: 11247548]
126. Luo J, Shen L & Zheng D Association between vitamin C intake and lung cancer: a dose-response meta-analysis. *Sci. Rep* 4, 6161 (2014). [PubMed: 25145261]
127. Harris HR, Orsini N & Wolk A Vitamin C and survival among women with breast cancer: a meta-analysis. *Eur. J. Cancer* 50, 1223–1231 (2014). [PubMed: 24613622]
128. Harris HR, Bergkvist L & Wolk A Vitamin C intake and breast cancer mortality in a cohort of Swedish women. *Br. J. Cancer* 109, 257–264 (2013). [PubMed: 23736027]
129. Wohlrab C, Phillips E & Dachs GU Vitamin C transporters in cancer: current understanding and gaps in knowledge. *Front. Oncol* 7, 74 (2017). [PubMed: 28484682]
130. Michels AJ, Hagen TM & Frei B Human genetic variation influences vitamin C homeostasis by altering vitamin C transport and antioxidant enzyme function. *Annu. Rev. Nutr* 33, 45–70 (2013). [PubMed: 23642198]
131. Savini I et al. SVCT1 and SVCT2: key proteins for vitamin C uptake. *Amino Acids* 34, 347–355 (2008). [PubMed: 17541511]
132. Wright ME et al. Genetic variation in sodium-dependent ascorbic acid transporters and risk of gastric cancer in Poland. *Eur. J. Cancer* 45, 1824–1830 (2009). [PubMed: 19243932]
133. Casabonne D et al. Fruit and vegetable intake and vitamin C transporter gene (SLC23A2) polymorphisms in chronic lymphocytic leukaemia. *Eur. J. Nutr* 56, 1123–1133 (2017). [PubMed: 26838684]
134. Duell EJ et al. Vitamin C transporter gene (SLC23A1 and SLC23A2) polymorphisms, plasma vitamin C levels, and gastric cancer risk in the EPIC cohort. *Genes Nutr* 8, 549–560 (2013). [PubMed: 23737080]
135. Mastrangelo D et al. Mechanisms of anti-cancer effects of ascorbate: Cytotoxic activity and epigenetic modulation. *Blood Cells. Mol. Dis* 69, 57–64 (2018). [PubMed: 28954710]
136. Bienert GP & Chaumont F Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochim. Biophys. Acta* 1840, 1596–1604 (2014). [PubMed: 24060746]
137. Uetaki M et al. Metabolomic alterations in human cancer cells by vitamin C-induced oxidative stress. *Sci. Rep* 5, 13896 (2015). [PubMed: 26350063]
138. Li R Vitamin C, a multi-tasking molecule, finds a molecular target in killing cancer cells. *React. Oxyg. Species* 1, 141–156 (2016).
139. Young JI, Züchner S & Wang G Regulation of the epigenome by vitamin C. *Annu. Rev. Nutr* 35, 545–564 (2015). [PubMed: 25974700]
140. Camarena V & Wang G The epigenetic role of vitamin C in health and disease. *Cell. Mol. Life Sci* 73, 1645–1658 (2016). [PubMed: 26846695]
141. Denko NC Hypoxia, HIF1 and glucose metabolism in the solid tumour. *Nat. Rev. Cancer* 8, 705–713 (2008). [PubMed: 19143055]
142. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT01852890> (2018).
143. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT01752491> (2018).
144. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03797443> (2019).
145. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03410030> (2019).
146. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT02420314> (2019).
147. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03613727> (2018).
148. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT02344355> (2018).

149. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03146962> (2018).
150. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT02905591> (2018).
151. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT02905578> (2018).
152. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT02516670> (2019).
153. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03334409> (2019).
154. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03418038> (2018).
155. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT03015675> (2017).
156. US National Library of Medicine. ClinicalTrials.gov <https://clinicaltrials.gov/ct2/show/NCT02969681> (2019).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Box 1 |**Transport of vitamin C**

The highest tissue concentrations of vitamin C are found in the brain, the adrenal gland and white blood cells, with concentrations ranging from 1 to 15 mM. These concentrations are 15–200 times higher than those in the plasma owing to active transport mechanisms⁴, mainly via sodium-dependent vitamin C transporters (SVCT1 and SVCT2)¹¹⁵. SVCT1 is mainly expressed in intestinal and renal epithelial cells, where it mediates absorption and re-absorption of vitamin C, respectively. SVCT2 is expressed throughout the entire body and is considered to be the primary vitamin C transporter. Interestingly, 50% of SVCT1-knockout mice do not survive until weaning, and the deletion of SVCT2 in mice leads to neonatal death^{116,117}.

Unlike ascorbate, dehydroascorbic acid (DHA) is transported by a class of facilitative glucose transporters (GLUTs)¹¹⁸. Among more than 12 different GLUTs, GLUT1 and GLUT3 have a higher affinity for DHA than for glucose⁶. However, under physiological conditions, GLUT transporters are unlikely the dominant path for ascorbate accumulation in most tissues, because glucose levels in the plasma (2–5 mM) are significantly higher than DHA levels (5–10 μ M). Red blood cells, and certain cancer cells, do not express SVCTs but transport vitamin C mainly as DHA via GLUT1 (REF.⁸). The rate of DHA uptake via GLUT1 or GLUT3 is at least 10–20 times faster than ascorbate uptake via SVCTs¹¹⁹. This difference is because the highly favourable reduction in intracellular DHA to ascorbate drives DHA uptake by cells. Furthermore, local DHA concentrations in body fluids can be higher under pathological conditions, such as cancer, where ROS released from cancer cells can facilitate the oxidation of extracellular ascorbate to DHA^{54,55,120}. For example, intestinal tumours harbouring *KRAS* mutations exhibited an increase in intracellular ascorbate level from 100 μ M (basal level) to more than 10 mM within 1 hour following intraperitoneal injection of high-dose ascorbate to a mouse model of intestinal tumours (*Apc*^{-/-}; *Kras*^{G12D+}) compared with tumours without *Kras* mutation (*Apc*^{-/-})⁴⁷. This drastic increase in intracellular ascorbate level in *KRAS* mutant tumours can be explained by the selective uptake of DHA via GLUT1 and its subsequent reduction to ascorbate. Tumours from *Apc*^{-/-}; *Kras*^{G12D+} mice highly express GLUT1 compared with *APC*^{-/-} mice, whereas both tumours express low levels of SVCTs.

Box 2 |**Oral intake of high-dose vitamin C and cancer prevention**

Patients with cancer often have lower plasma concentrations of ascorbate than healthy adults^{121,122} and vitamin C deficiency is associated with an increased risk of cancer mortality^{123,124}. In Western countries, plasma ascorbate levels can vary among individuals, but the lowest quartile of men have a significantly higher risk of mortality from cancer¹²⁵. In a meta-analysis of 21 studies, including ~9,000 lung cancer cases¹²⁶, a correlation was shown between an individual's risk of lung cancer and their vitamin C intake, where male adults in the USA who took 100 mg per day of dietary vitamin C had a 7% reduced lung cancer risk. This dose is also associated with a reduced overall mortality and breast-cancer-specific mortality in women^{127,128}. That vitamin C may prevent or delay cancer development was further supported by preclinical studies. In ascorbate-deficient *Gulo*^{-/-} mice, ten-eleven translocation 2 (TET2) deletion drives development of acute myelogenous leukaemia (AML)⁷², which is suppressed by administration of oral ascorbate. Similarly, administration of oral ascorbate 7 days before inoculation of cancer cells decreased tumour development in a lymphoma xenograft model⁷⁹. Taken together, oral doses of vitamin C may be an effective agent to prevent the development of certain types of malignancy, especially for individuals who may have mutations such as TETs predisposing them to cancer.

Genetic variation in ascorbate transporters has also been found to associate with cancer risk¹²⁹⁻¹³¹. Although the correlation with cancer remains weak and plasma ascorbate was often not measured, single-nucleotide polymorphisms (SNPs) in the sodium-dependent vitamin C transporters *SLC23A1* (SVCT1) and *SLC23A2* (SVCT2) have been linked to the risk of certain cancers^{132,133}. Some of these SNPs are associated with reduced levels of ascorbate systemically, which may have adverse consequences for ascorbate-dependent biochemical processes that involve, for example, Fe²⁺-dependent and α -ketoglutarate-dependent dioxygenases¹³⁴. However, it remains unclear whether the reduction in ascorbate levels associated with those SNPs can be directly attributed to compromised function of transporters. Together, these findings support the profound importance of the amount of oral ascorbate to humans' optimized health and disease prevention. However, to address the definite benefits of supplementary oral intake of ascorbate, potentially in combination with therapeutic intravenous ascorbate therapy, rigorous preclinical and randomized clinical trials will be required.




Fig. 1. Integrated pro-oxidant mechanism of vitamin C and cancer cell cytotoxicity,
a | Ascorbate can be oxidized in the extracellular space by reactive oxygen species (ROS), producing ascorbate radical, which can be oxidized to dehydroascorbic acid (DHA). DHA can be taken up by cells or irreversibly converted to 2,3-L-diketoglutonate (2,3-DKG), which is degraded into oxalic acid and threonic acid, **b** | Pharmacological ascorbate can kill cancer cells by increasing oxidative stress via two possible mechanisms that complement each other. First, extracellular H₂O₂ may directly kill cancer cells by generating •OH via the Fenton reaction^{25,26,135}. Increased levels of labile ferric iron, Fe³⁺, in the tumour microenvironment can facilitate the oxidation of ascorbate, resulting in ascorbate radical, DHA and ferrous iron, Fe²⁺. Once Fe²⁺ is formed, Fe²⁺ may be oxidized by oxygen, producing superoxide anions, O₂•⁻. Superoxide dismutase (SOD) catalyses the conversion of O₂•⁻ to H₂O₂ and O₂. Fe³⁺ can enter the cell when bound to transferrin (Tf), which binds to the Tf receptor (TfR) and is processed and oxidized in the endosome to then contribute to the intracellular Fe²⁺ pool³⁶. H₂O₂ can enter the cell through diffusion facilitated by aquaporins¹³⁶. H₂O₂ reacts with either extracellular or intracellular labile Fe²⁺ to generate

highly reactive hydroxyl radicals ($\cdot\text{OH}$) that are harmful to cells. These reactions are further perpetuated by the recycling of Fe^{3+} to Fe^{2+} by ascorbate and ascorbate radical, generating fully oxidized vitamin C, DHA. Second, H_2O_2 may contribute to the increased levels of extracellular DHA by creating a more oxidative tumour microenvironment. DHA can then efficiently enter cells through glucose transporter 1 (GLUT1) and consume the intracellular reducing potential of reduced glutathione (GSH) and NADPH, resulting in increased levels of intracellular ROS^{51,52}. This leads to poly(ADP-ribose) polymerase (PARP) activation, a DNA repair enzyme, thereby depleting cellular NAD^+ levels, a cofactor of PARP. NAD^+ is required by glyceraldehyde 3-phosphate dehydrogenase (GADPH) as a cofactor. Consequent inhibition of GAPDH activity inhibits glycolysis in cancer cells, leading to inhibition of ATP production and cell death^{47,137,138}. In addition, cellular ROS can also be released from cells, resulting in a positive feedback loop. Because high levels of labile Fe^{2+} , GLUT1 overexpression and addiction to glycolysis frequently occur in many types of cancer cells, certain cancer cells may present all three of these characteristics and those populations might be more sensitive to ascorbate treatment. 1,3BPG, 1,3-bisphosphoglyceric acid; G3P, glyceraldehyde 3-phosphate; G6PD glucose-6-phosphate dehydrogenase; GSSG, glutathione disulfide; PPP, pentose phosphate pathway; SVCT, sodium-dependent vitamin C transporters.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 2 |. Regulation of TET enzymes by ascorbate.

Catalysed by DNA methyltransferases (DNMTs), DNA methylation occurs at the carbon-5 position of cytosine. Intracellular ascorbate influences the DNA methylation landscape by enhancing the enzymatic activity of ten-eleven translocation enzymes (TETs), which actively remove cytosine methylation marks through a series of oxidation reactions dependent on oxygen, α-ketoglutarate (αKG), Fe²⁺ and ascorbate on the basis of its function as an αKG-dependent dioxygenase (αKGDD)^{139,140}. TETs first convert 5-methylcytosine (5mC) to 5-hydroxymethyl-cytosine (5hmC). In the next two steps, 5hmC is further oxidized to 5-formylcytosine (5fC) and 5-carboxylcytosine (5CaC). Subsequently, 5fC and 5CaC are converted to cytosine by the base excision repair pathway enzyme, thymine DNA glycosylase. By promoting the recycling of Fe³⁺ to Fe²⁺, ascorbate ensures that TETs are constantly active. DHA, dehydroascorbic acid.




Fig. 3 | Ascorbate and HIF1 α regulation.

Ascorbate is a vital cofactor for the hypoxia-inducible factor (HIF) hydroxylases, proline hydroxylase domain proteins (PHDs) and asparagine hydroxylase (factor-inhibiting HIF (FIH)), which are also members of the α -ketoglutarate (α KG)-dependent dioxygenase (α KGDD) protein family^{77,141}. HIF1, a heterodimeric transcription factor, consists of two sub-units: HIF1 α , regulated by O₂, and HIF1 β . Under normal conditions with sufficient oxygen and ascorbate availability, the functional capacity of HIF1 α is inhibited by the HIF hydroxylases. HIF1 α is hydroxylated at proline residues by PHD. Prolyl-hydroxylated HIF1 α is then bound by the von Hippel-Lindau (VHL) tumour suppressor protein, which recruits an E3-ubiquitin ligase that targets HIF1 α for proteasomal degradation and thus limits the quantity of HIF1 α units within the cell. HIF1 α activity is regulated within the nucleus and can be inhibited. FIH hydroxylates an asparagine residue, N806, on HIF1 α . This hydroxyl group prevents p300, a co-activator protein, from associating with the HIF complex resulting in the inhibition of the transcription activity of HIF1 and the activation of any downstream pathways. High-dose ascorbate treatment in tumour tissues with normoxic HIF1 α stabilization can potentially increase PHD and FIH activity to degrade HIF1 protein and slow down tumour growth. In conditions where ascorbate is depleted, such as in certain cancer types or in tumour tissue in *Gulo*^{-/-} mice, the activity of PHD and FIH is reduced even when oxygen is available, which leads to stabilization and activation of HIF1 α and its translocation to the nucleus. HIF1 α associates with HIF1 β , p300 and other cofactors within the nucleus to induce target genes such as *GLUT1*, which together might promote tumour growth.

Table 1 I

List of active intravenous vitamin C clinical trials as an anticancer therapy

Phase	Trial name	Institution	Trial design	Cancer type	Drugs	Refs
Phase I	Gemcitabine, ascorbate, radiation therapy for pancreatic cancer	Holden Comprehensive Cancer Center at the University of Iowa (USA)	Single arm	Pancreatic neoplasms	Gemcitabine; radiation + IV ascorbate	142
Phase I	High-dose ascorbate in glioblastoma multiforme	Holden Comprehensive Cancer Center at the University of Iowa (USA)	Single arm	Glioblastoma	Radiation + temozolomide + IV ascorbate	143
Phase I/II	High-dose ascorbate + nanoparticle paclitaxel protein bound + cisplatin + gemcitabine in patients who have had no prior therapy for their metastatic pancreatic cancer	Piedmont Cancer Institute (USA)	Single arm	Metastatic pancreatic cancer	Paclitaxel, cisplatin, gemcitabine + IV ascorbate	144
Phase I/II	High-dose ascorbate+ nanoparticle paclitaxel protein bound + cisplatin + gemcitabine in patients who have no prior therapy for their metastatic pancreatic cancer	HonorHealth Research Institute, University of California-San Diego Moores Cancer Center and Piedmont Cancer Institute (USA)	Single arm	Pancreatic cancer	Paclitaxel protein-bound, cisplatin + IV ascorbate	145
Phase II	High-dose ascorbate in stage IV non-small-cell lung cancer	Holden Comprehensive Cancer Center at the University of Iowa (USA)	Single arm	Non-small-cell lung cancer	Paclitaxel, carboplatin + IV ascorbate	146
Phase II	Therapeutic use of IV vitamin C in allogeneic stem cell transplant recipients	Virginia Commonwealth University-Massey Cancer Center (USA)	Single arm	Hodgkin lymphoma, lymphoid leukaemia and multiple myeloma	IV and oral vitamin C	147
Phase II	Pharmacological ascorbate combined with radiation and temozolomide in glioblastoma multiforme: a phase II trial	Holden Comprehensive Cancer Center at the University of Iowa (USA)	Single arm	Glioblastoma multiforme	Radiation + temozolomide + IV ascorbate	148
Phase II	High-dose vitamin C IV infusion in patients with resectable metastatic solid tumor malignancies	Weill Cornell Medicine (USA)	Single arm	Colorectal, pancreatic and lung cancer with <i>KRAS</i> or <i>BRAF</i> mutation	IV ascorbate	149
Phase II	Pharmacological ascorbate with concurrent chemotherapy and radiation therapy for non-small-cell lung cancer	Holden Comprehensive Cancer Center at the University of Iowa (USA)	Single arm	Non-small-cell lung cancer	Radiation, paclitaxel, carboplatin + IV ascorbate	150
Phase II	Pharmacological ascorbate, gemcitabine, nab-paclitaxel for metastatic pancreatic cancer	Holden Comprehensive Cancer Center at the University of Iowa (USA)	Randomized two-arm	Pancreatic neoplasms	Paclitaxel, gemcitabine + IV ascorbate	151
Phase II	Ascorbic acid in combination with docetaxel in men with metastatic prostate cancer	Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins (USA)	Randomized two-arm	Hormone-resistant prostate cancer, metastatic prostate carcinoma and stage IV prostate cancer	IV ascorbate+ docetaxel	152
Phase II	IV ascorbic acid as an adjunct to pazopanib in the first-line setting for metastatic or unresectable clear cell renal cell carcinoma (ccRCC)	Mayo Clinic in Florida and Minnesota, Illinois CancerCare-Peoria, Iowa-Wide Oncology Research Coalition NCCORP and	Randomized two-arm	Clear cell renal cell carcinoma	Pazopanib hydrochloride + IV ascorbate	153

Phase	Trial name	Institution	Trial design	Cancer type	Drugs	Refs
Phase II	Ascorbic acid and combination chemotherapy in treating patients with relapsed or refractory lymphoma	Sanford Medical Center Fargo (USA) Mayo Clinic in Arizona, Minnesota and Florida and Holden Comprehensive Cancer Center at the University of Iowa (USA)	Randomized two-arm	B cell lymphoma with MYC and BCL2/BCL6 rearrangement, and recurrent Hodgkin lymphoma	Carboplatin, cisplatin+IV ascorbate	154
Phase III	IV ascorbic acid in advanced gastric cancer	Sun Yat-Sen University Cancer Center (China)	Randomized two-arm	Gastric cancer	mFOLFOX6 + IV ascorbate	155
Phase III	IV ascorbic acid in combination with FOLFOX +/- bevacizumab versus treatment with FOLFOX +/- bevacizumab alone as first-line therapy for advanced colorectal cancer	Sun Yat-Sen University Cancer Center (China)	Randomized two-arm	Colorectal neoplasms	mFOLFOX6, bevacizumab + IV ascorbate	156

IV, intravenous.